

Mini-math Div 3/4: Friday, October 25, 2024 (7.1-7.5) - 18 minutes

SOLUTIONS

1. Write a differential equation that describes the following relationships. If necessary, use k as the constant of proportionality.

- (a) (2 points) The rate of change of population, P , with respect to time, t , is inversely proportional to the square root of time and directly proportional to the area, A , that the population covers.

Solution:

$$\frac{dP}{dt} = \frac{kA}{\sqrt{t}}$$

- (b) (2 points) The position of a particle is given by $s(t)$, where t is measured in seconds. Its acceleration is directly proportional to its position. When the particle is at position 4 units, its acceleration is 2 units/ s^2 .

Solution:

$$\frac{d^2s}{dt^2} = \frac{1}{2}s$$

2. (4 points) Determine the value of k , if any, for which $y = \sin(2x) - k \sin(4x)$ would be a solution to the differential equation $y'' + 4y = 3 \sin(4x)$.

Solution:

$$y' = 2 \cos(2x) - 4k \cos(4x)$$

$$y'' = -4 \sin(2x) + 16k \sin(4x)$$


Then

$$[-4 \sin(2x) + 16k \sin(4x)] + 4 [\sin(2x) - k \sin(4x)] = 3 \sin(4x)$$

so $16k - 4k = 3$, giving $k = 1/4$.

3. (2 points) What differential equation can the slope field to the right represent?

- A. $\frac{dy}{dx} = -x/y$
- B. $\frac{dy}{dx} = -y/x$
- C. $\frac{dy}{dx} = y^2$
- D. $\frac{dy}{dx} = x/y$
- E. $\frac{dy}{dx} = y/x$




Solution:

At $x = 0$, the slopes are 0, so (A) or (D). Since the slopes are positive for $x > 0$ and $y < 0$, (D) is the answer.

4. (2 points) The slope field for a certain differential equation is shown to the right. Which of the following could be a particular solution to the differential equation?

- A. $y = x^3$
- B. $y = \frac{1}{x+2}$
- C. $y = -2^x - 2$
- D. $y = e^{-x} - 2$
- E. $y = e^x + 2$



Solution: The solutions have a horizontal asymptote as $x \rightarrow -\infty$, so (C) is the answer.

5. Consider the initial value problem $\frac{dy}{dx} = 2x + y$ and $y(1) = 2$.

(a) (2 points) Find an approximation of $y(1.2)$ using Euler's Method with two equal steps.

Solution:

$$y(1.1) \approx y(1) + y'(1)(1.1 - 1) = 2 + (2(1) + 2)(0.1) = 2.4,$$

$$y(1.2) \approx y(1.1) + y'(1.1)(1.2 - 1.1) \approx 2.4 + (2(1.1) + 2.4)(1.2 - 1.1) = 2.4 + (4.6)(0.1) = 2.86$$

(b) (2 points) Is your estimate in part (a) an overestimate or an underestimate?

Solution: At $(1, 2)$,

$$\frac{d^2y}{dx^2} = 2 + \frac{dy}{dx} = 2 + 2x + y > 0$$

so the function is concave up. Therefore, the estimate will be an underestimate.

(Aside: the actual value is about 2.92842. If we used 10 equal steps, we would get the approximation 2.91397)