Mini-math Div 3/4: Wednesday, October 30, 2024 (7.6-7.9) (20 minutes) SOLUTIONS

1. (2 points) Solve the following differential equation:

$$\frac{dy}{dx} = xy\sin(x^2) \cdot \ln y$$

Solution:

$$\int \frac{1}{y \ln y} \, dy = \int x \sin x^2 \, dx$$
$$\ln |\ln y| = -\frac{1}{2} \cos x^2 + C_1$$
$$|\ln y| = e^{-\frac{1}{2} \cos x^2 + C_1}$$
$$\ln y = \pm e^{-\frac{1}{2} \cos x^2 + C_1} \quad \text{or} \quad C_2 e^{-\frac{1}{2} \cos x^2}$$
$$y = e^{\pm e^{-\frac{1}{2} \cos x^2 + C_1}} \quad \text{or} \quad e^{C_2 e^{-\frac{1}{2} \cos x^2}}$$

- 2. (2 points) During a chemical reaction, the rate of change of the amount of the chemical remaining is proportional to the amount remaining. At time t = 0, the amount of the chemical is 60 g. At time t = 8, the amount of the chemical is 12 g. At what time t is the amount of the chemical 4 g?
 - A. $\frac{4\sqrt{42}}{3}$ B. $\frac{28}{3}$ C. $\frac{8\ln 15}{\ln 5}$ D. $\frac{8\ln 4}{\ln 12}$ E. $\frac{8}{3}$

Solution:

$$P(t) = P_0 e^{kt} = P_0 (e^k)^t$$

$$P_1 = P_0 (e^k)^{t_1} \implies e^k = \left(\frac{P_1}{P_0}\right)^{1/t_1}$$

$$P(t) = P_0 \left(\frac{P_1}{P_0}\right)^{t/t_1}$$

$$P_2 = P_0 \left(\frac{P_1}{P_0}\right)^{t_2/t_1} \implies t_2 = \frac{t_1 \ln(P_2/P_0)}{\ln(P_1/P_0)} = \frac{8 \ln(1/15)}{\ln(1/5)} = \frac{8 \ln 15}{\ln 5}$$
(C)

3. (2 points) Solve the following initial value problem:

$$\frac{dy}{dx} = y^2, \quad y(3) = -2$$
A. $y = \frac{1}{5/2 - x}$ for $x \neq 5/2$
B. $y = \frac{2}{5 - 2x}$ for $x > 5/2$
C. $y = -\frac{1}{x} - \frac{5}{3}$ for $x \neq 0$
D. $y = -\frac{5x + 3}{3x}$ for $x > 0$

Solution:

$$\int \frac{1}{y^2} dy = \int 1 dx \quad \Rightarrow \quad -\frac{1}{y} = x + C,$$
$$\frac{1}{2} = 3 + C \quad \Rightarrow C = -\frac{5}{2},$$
$$y = \frac{1}{5/2 - x} = \frac{2}{5 - 2x}$$

Since the domain is the largest open interval which contains the initial condition, the domain is x > 5/2.

(B)

- 4. The number of squirrels in a park at time t is modelled by the function y = F(t) that satisfies the logistic differential equation $\frac{dy}{dt} = \frac{1}{2000}y(1500 - y)$, where $t \ge 0$ is measured in weeks. The number of squirrels in the park at time t = 0 is F(0) = b, where b is a positive constant.
 - (a) i. (1 point) If b = 300, what is the largest rate of increase in the number of squirrels in the park?

Solution: Since 300 < 1500/2 = 750, F grows most rapidly when it is half the carrying capacity, 750. Then

$$\left. \frac{dy}{dt} \right|_{F=750} = \frac{750}{2000} (1500 - 750) = \frac{1125}{4} = 281.25$$

ii. (1 point) If b = 1000, what is the largest rate of increase in the number of squirrels in the park?

Solution: For F > 750, $\frac{dy}{dt} > 0$ and $\frac{d^2y}{dt^2} < 0$, so F grows most rapidly when F = 1000.

$$\left. \frac{dy}{dt} \right|_{F=1000} = \frac{1000}{2000} (1500 - 1000) = 250$$

(b) (2 points) If b = 150, find $\lim_{t \to \infty} F(t)$ and interpret the meaning of this limit in the context of the problem. For reference, the differential equation is $\frac{dy}{dt} = \frac{1}{2000}y(1500 - y)$.

Solution: The carrying capacity is 1500, so $\lim_{t\to\infty} F(t) = 1500$. This means in the long term, the population of squirrels in the park will tend to 1500.

(c) (4 points) (*) Find the function F(t) if b = 500.

Solution:

$$\begin{aligned} \int \frac{dy}{y(1500-y)} &= \int \frac{dt}{2000} \\ \int \left(\frac{1/1500}{y} + \frac{1/1500}{1500-y}\right) &= \int \frac{dt}{2000} \\ \ln \left|\frac{y}{1500-y}\right| &= (\ln |y| - \ln |1500-y|) = \frac{1500}{2000}x + C = \frac{3}{4}x + C \\ \frac{y}{1500-y} &= Ce^{3t/4} \end{aligned}$$

Using the initial condition,

$$\frac{1}{2} = \frac{500}{1500 - 500} = C$$

 \mathbf{SO}

$$\frac{2y}{1500 - y} = e^{3t/4}$$
$$2y = 1500e^{3t/4} - e^{3t/4}y$$
$$y = \frac{1500e^{3t/4}}{e^{3t/4} + 2}$$

(Alternatively, go directly to the general solution if you have memorized it. Be careful with using the correct general solution which depends on the form of the logistic DE.)