

Mini-math Div 3/4: Friday, March 19, 2025 (10.10-10.15) - (25 minutes)

SOLUTIONS

1. (3 points) The series  $\sum_{n=1}^{\infty} \frac{(-1)^n}{n}$  converges to  $S$ . If  $S_n$  is used to approximate  $S$ , what is the least value of  $n$  for which the alternating series error bound guarantees an error to strictly within 0.01?

**Solution:** Notice the series is alternating, and  $1/n$  is decreasing and limits to 0. Then by the Alternating Series Test error bound,

$$\begin{aligned} \frac{1}{n+1} &< \frac{1}{100} \\ n+1 &> 100 \end{aligned}$$

so  $n = 100$  is the least value of  $n$  needed.

2. (3 points) Let  $P(x)$  be the fifth-degree Taylor Polynomial for a function  $f$  about  $x = 1$ . Information about the maximum of the absolute value of selected derivatives of  $f$  over various intervals is given below.

$$\begin{aligned} \max_{0 \leq x \leq 1.5} |f^{(4)}(x)| &= 4.6, & \max_{0 \leq x \leq 1.5} |f^{(5)}(x)| &= 7.2, & \max_{0 \leq x \leq 1.5} |f^{(6)}(x)| &= 6.8, \\ \max_{1 \leq x \leq 1.5} |f^{(4)}(x)| &= 3.2, & \max_{1 \leq x \leq 1.5} |f^{(5)}(x)| &= 4.7, & \max_{1 \leq x \leq 1.5} |f^{(6)}(x)| &= 5.1 \end{aligned}$$

Find the smallest value of  $k$  for which the Lagrange error bound guarantees that

$$|f(1.5) - P(1.5)| \leq k$$

**Solution:** We use the bound on the 6th derivative on the interval  $[1, 1.5]$ , since this is the smallest interval which contains the centre and the point we are approximating. The Lagrange error bound gives a maximum absolute error of

$$\frac{\max_{1 \leq x \leq 1.5} |f^{(6)}(x)|}{6!} \cdot |1.5 - 1|^6 = \frac{5.1 \cdot 0.5^6}{6!}$$

3. (4 points) Find the interval of convergence for the series  $\sum_{n=1}^{\infty} \frac{(-1)^n (x-3)^n}{n2^n}$

**Solution:**

$$\lim_{n \rightarrow \infty} \left| \frac{(-1)^{n+1} (x-3)^{n+1}}{(n+1)2^{n+1}} \cdot \frac{n2^n}{(-1)^n (x-3)^n} \right| = \lim_{n \rightarrow \infty} \frac{n}{n+1} \cdot \frac{|x-3|}{2} = \frac{|x-3|}{2}$$

Then the radius of convergence is 2. Testing  $|x-3|=2$ , we have at  $x-3=2$

$$\sum_{n=1}^{\infty} \frac{(-1)^n 2^n}{n2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$$

converges by AST and

$$\sum_{n=1}^{\infty} \frac{(-1)^n (-2)^n}{n2^n} = \sum_{n=1}^{\infty} \frac{1}{n}$$

diverges by  $p$ -series. Therefore, the interval of convergence is  $(3-2, 3+2] = (1, 5]$ .

4. (3 points) What is the Maclaurin series for  $\frac{\cos x - 1}{x}$ ? Assume differentiability at 0 (e.g. the function has a value at 0 which makes it differentiable). You may, but are not required to, express your answer in summation notation.

**Solution:**

$$\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \dots$$

so

$$\begin{aligned} \frac{\cos x - 1}{x} &= \frac{1}{x} \cdot \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} = \frac{-\frac{x^2}{2} + \frac{x^4}{4!} - \dots}{x} \\ &= \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n-1}}{(2n)!} = -\frac{x}{2} + \frac{x^3}{4!} - \dots \end{aligned}$$

5. (4 points) Let  $f$  be a function with  $f(0) = 2$  and  $f'(x) = \arctan x$ . Write the first three non-zero terms of the Maclaurin series for  $f$ .

**Solution:**

$$\begin{aligned}\arctan x &= x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \\ f(x) &= \int \arctan x \, dx = \int \left( x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \right) dx \\ &= C + \frac{x^2}{2} - \frac{x^4}{4 \cdot 3} + \frac{x^6}{6 \cdot 5} - \dots\end{aligned}$$

Since  $f(0) = 2$ , we get  $C = 2$  so the first three non-zero terms are:

$$2 + \frac{x^2}{2} - \frac{x^4}{12}$$